



Foetal Alcohol Syndrome / Alcohol Related Neurodevelopmental Disorder

First description and alternative names

FAS was first observed in Nantes by paediatrician Paul Lemoine in 1968 who described similar dysmorphic facial features and growth delays in 127 infants of pregnant alcohol-drinking mothers. Fetal Alcohol Syndrome (FAS) was coined in Seattle by David Smith and Ken Jones in 1973(1,2). The Institute of Medicine in 1996 delineated Full FAS, Partial FAS, and Alcohol Related Neurodevelopmental Disorder (ARND) based on the presence or absence of facial dysmorphism and /or growth retardation (3). ARND replaced the old term Fetal Alcohol Effect (FAE). FAS Spectrum Disorders, an umbrella term, was introduced by O'Malley & Hagerman in 1998, refined to Fetal Alcohol Spectrum Disorders (FASDs) by Streissguth & O'Malley in 2000 (4,5). In 2013 DSMV proposed a new diagnostic guidelines for those with neurodevelopmental disorders associated with prenatal alcohol exposure (NDDPAE 315.8) but without facial features. It requires features to be ruled into a diagnosis with other factors ruled out. This was the first time this was included in an international diagnostic manual.

Genetics and molecular biology

Both FAS and ARND are teratogenic conditions, but recent research is beginning to study the epigenetic effects of prenatal alcohol exposure. For example, the major liver pathway of oxidative metabolism which produces acetaldehyde by cytotoxic alcohol dehydrogenase is also accompanied by a reduction in NAD to NADH. There is a subsequent alteration in the cellular redox triggered by a change in the NAD/NADH ratio which may result in gene activation resulting in a change in gene expression. Increasingly the impact of prenatal alcohol on epigenetic mechanisms has also been investigated. For example studies have demonstrated that prenatal alcohol exposure has the ability to modify methylation of the retrotransposon prior to the AVY gene in genetically inbred mice, leading to differences in coat colors (17). A wide range of mechanisms beyond this have been identified, from direct apoptotic damage, interneuronal signaling deficits and damage to scaffolding proteins interfering with neural migration (18).

Incidence/ prevalence

The recognition and incidence for FAS has been shown to be increasing. Original estimates varied from 3 per 1,000 live births in the USA to 40 per 1,000 births in South Africa. More recent studies in Lazio, Italy and Cape Town, South Africa, have suggested much higher rates of 35/1000 to 890/1000 respectively (6,10,13). Methodological issues, genetic differences in the susceptibility to alcohol based on the mother's liver metabolism, as well as differences in population drinking patterns may account for some of the variance(7). As a result, whilst the exact incidence for any one population differs greatly and is unknown the rates are considered potentially higher than previously thought. No documentation of decreased life expectancy exists.

Physical features and psychiatric characteristics

Characteristic facial features include short palpebral fissures, thin (flat) upper lip, flattened philtrum, flat midface. Growth retardation less than the 10th percentile for height and weight. Low birth weight for gestational age, decelerating weight over time not due to nutrition, disproportional low weight-to-height ratio. FAS has the classic facial features, ARND does not

have the facial features. Gross and fine motor delays as in Developmental Coordination Disorder. Alcohol Related Birth Defects (ARBD) with eye, renal, cardiac and/or skeletal anomalies. FAS is the much less common, but most recognisable form of FASD (3,8,9,10).

Infants can demonstrate Primary Regulatory Disorders characterized by difficulties in tolerating environmental stimuli (i.e. habituation problems), co-ordinating basic motor movements (i.e. sucking), or interacting with parents or care-providers. Secondary psychiatric disorders appear due to environmental stressors such as PTSD after physical or sexual abuse, or Reactive Attachment Disorder of Infancy or Early Childhood related to separation from birth mother or multiple foster home placements. FASD increases the vulnerability to many common psychiatric disorders, such as ADHD, Mood Disorder, Anxiety Disorder, Alcohol or Drug Dependence, PDD, Autism and Schizoaffective Disorder. Personality Disorders seen are, Avoidant, Dependent, Schizoid, Passive/Aggressive and Borderline. Presence or absence of facial dysmorphology or growth features do not clinically correlate with the psychiatric presentation or structural brain damage in FASD (5,8, 11, and 12).

Neuropsychological Deficits

70-75% of patients with FASD do not have intellectual disability. The neuropsychological profile consistently shows a Complex Verbal and Non-Verbal Learning Disorder affecting multiple domains of functioning including attention, impulsivity, working memory, executive function, processing speed, social skills, interpersonal relatedness and language development. It includes a Mathematics Disorder and/ or Disorder of Written Expression and/or Reading Disorder, and evidence of a Mixed Receptive/ Expressive Language Disorder with specific deficits in social cognition and social communication. Vineland Adaptive Behavioral Scale (VABS) shows Functional Deficits in daily living skills, socialization and communication. Those with higher functioning in some areas can often mask their difficulties until external pressures lead to higher level abilities such as executive functioning being less effective. Simple functions are often intact. For example an individual can sequence and switch separately but not when these two tasks are combined. Working memory deficits tend to be verbal working memory deficits rather than numerical having implication as to how these skills are tested. (3, 5, 8,9,10, 13).

Brain structural abnormalities

Autopsy reports of infant deaths and MRI studies showed diffuse structural brain changes. These include hydrocephalus, microcephaly, corpus callosal agenesis, cerebellar hypoplasia, enlarged ventricles and heterotopias (8, 9). Brain imaging studies consistently show microcephaly. The corpus callosum is most commonly affected. The basal ganglion caudate and the hippocampus, integral in working memory and executive function, may be decreased in shape and volume, and the cerebellum may be smaller (5,9,14).

Brain neurotransmitter and neurophysiological abnormalities

Diffuse CNS neurotoxic effects include cell death and/or disruption of cell migration, proliferation, differentiation and maturation (3,5, 8, and 9). Deficits have been discovered in all neurotransmitter systems. Dopaminergic and noradrenergic deficits are likely connected to ADHD presentation of FASD (4,12,15). EEG abnormalities show infant/ child cerebral immaturity, sleep state EEG, as well as temporal lobe dysrhythmia and complex partial seizure disorder in 6 to 19 year olds with FASD (1,12).

Available guidelines for behavioral assessment/ treatment/management strategies

Despite the high prevalence of the disorder, there have been few actual randomised studies of behavioural interventions specific to a FASD population. Those that have been have been small scale. Whilst some benefits are noted much more work is required. A review by Chandresena recently looked at possible strategies including medical, educational and psychological (16).

Useful websites /associations for more information

- www.fasdaware.co.uk
- www.fasdtrust.co.uk
- www.nofas.co.uk
- www.fasd.ie
- www.nofas.com

References

1. Lemoine P., Harousseau H., & Borteyru J.P. (1968) Les enfants de parents alcooliques: Anomalies observees a propos de 127 cas. *Quest Med* **21**, 476-482.
2. Jones K.L., & Smith D.W. (1973) Recognition of the fetal alcohol syndrome in early infancy. *Lancet* **2**, 999-1101.
3. Stratton K.R., Rowe C.J., & Battaglia F.C. (1996) *Fetal Alcohol Syndrome: Diagnosis, epidemiology, prevention and treatment in medicine*. Washington DC: National Academy Press.
4. O'Malley K.D., & Hagerman R.J. (1998) Developing Clinical Practice Guidelines for Pharmacological Interventions with Alcohol-affected Children. In: *Proceedings of a special focus session of the Interagency Co-ordinating Committee on Fetal Alcohol Syndrome*. Chevy Chase Ma, Sept 10th & 11th, (eds CDC & NIAAA), 145-177.
5. Streissguth A.P. & O'Malley K.D. (2000) Neuropsychiatric implications and long-term consequences of fetal alcohol spectrum disorders. *Semin Clin Neuropsychiat* **5**, 177-190.
6. May P.A., Gossage J.P., Kalberg W.O., Robinson L.K., Buckley D., Manning M., & Hoyme H.E. (2009) Prevalence and epidemiological characteristics of FASD from various research methods with an emphasis on in school studies. *Dev Disabil Res Rev* **15**, 176-192.
7. McCarver D.G., Thomasson H.R., Martier S.S., Sokol R.J., & Li T.K. (1997) Alcohol dehydrogenase*3 allele protects against alcohol-related birth defects among African Americans. *J Pharmacol Exp Ther* **283(3)**, 1095-1101.
8. Streissguth A.P. (1997) *Fetal alcohol syndrome. A guide for families and communities*. Baltimore:Brookes Publishing.
9. Hagerman R.J. (1999) Fetal Alcohol Syndrome. In: *Neurodevelopmental Disorders. Diagnosis and Treatment*. pp.3-59, New York, Oxford: Oxford University Press.
10. Chudley A.E., Conroy J., Cook J.L., Looock C., Rosales T., & LeBlanc N. (2005) Fetal alcohol spectrum disorder: Canadian guidelines for diagnosis. *Can Med Assoc J* **172(5 suppl)**, S1-21.
11. Famy C., Streissguth A.P., & Unis A.S. (1998) Mental illness in adults with fetal alcohol syndrome or fetal alcohol effects. *Am J Psychiat* **155(4)**, 552-554.
12. O'Malley K.D. & Storz L. (2003) Fetal alcohol spectrum disorder and ADHD: diagnostic implications and therapeutic consequences. *Expert Rev Neurotherapeutics*, July, **3 (4)**, 477-489.
13. Streissguth A.P., Bookstein F.L., Barr H.M., Sampson P.D., O'Malley K.D., & Young K. (2004) Risk factors for adverse risk outcomes in fetal alcohol syndrome and fetal alcohol effects. *Dev Behav Pediat* **25 (4)**, 228-238.
14. Bookstein F.L., Sampson P.D., Streissguth A.P., & Connor P.L. (2001) Geometric morphometrics of corpus callosum and subcortical structures in fetal alcohol affected brain. *Teratology* **64**, 4-32.

15. Sobrain S.K., Jones B.L., James H., Kamara F.N., & Holson R.R. (2005) Prenatal alcohol preferentially enhances reactivity of the dopamine D1 but not D2 or D3 receptors in offspring. *Neurotoxicol Teratol* **27**, 73-93.
16. Chandrasena A.N., Mukherjee R.A.S., & Turk J.T. (2009) Fetal Alcohol Spectrum Disorders: An Overview of Interventions for Affected Individuals. *Child Adol Ment Health* **14**(4), pp 162-167.
17. HAYCOCK, P.C. (2009) Fetal alcohol spectrum disorders: The epigenetic perspective. *Biology of Reproduction* **81**:607–617,
- 18 BMA Board of Science. *Fetal Alcohol Spectrum Disorders, a guide for healthcare practitioners*. London: BMA Publishing; 2007

Raja Mukharjee, Kieran D O'Malley, , May 2015

Copyright © 2015 R. Mukherjee, K. O'Malley

The SSBP hopes that readers will find the syndrome information sheets useful. They represent the views of the authors who kindly prepared them, and not necessarily those of the SSBP.